

# Regulatory Mimicry in *Listeria monocytogenes* Actin-Based Motility

Ryan Chong,<sup>1,3</sup> Rachel Swiss,<sup>1,3</sup> Gabriel Briones,<sup>1</sup> Kathryn L. Stone,<sup>2</sup> Erol E. Gulcicek,<sup>2</sup> and Hervé Agaisse<sup>1,\*</sup>

<sup>1</sup>Section of Microbial Pathogenesis, Boyer Center for Molecular Medicine, Yale University School of Medicine, New Haven, CT, USA

<sup>2</sup>Northeast Biodefense Center Proteomics Core, W.M. Keck Foundation Biotechnology Laboratory, New Haven, CT, USA

<sup>3</sup>These authors contributed equally to this work

\*Correspondence: [herve.agaisse@yale.edu](mailto:herve.agaisse@yale.edu)

DOI 10.1016/j.chom.2009.08.006

## SUMMARY

The actin-based motility of the intracellular pathogen *Listeria monocytogenes* relies on ActA, a bacterial factor with a structural domain allowing it to mimic the actin nucleation-promoting activity of host cell proteins of the WASP/WAVE family. Here, we used an RNAi-based genetic approach in combination with computer-assisted image analysis to investigate the role of host factors in *L. monocytogenes* cell-to-cell spread. We showed that the host cell serine/threonine kinase CK2 is required for efficient actin tail formation by *L. monocytogenes*. Furthermore, CK2-mediated phosphorylation of ActA regulated its affinity for the actin-nucleating ARP2/3 complex, as is the case for CK2-mediated phosphorylation of WASP and WAVE. Thus, ActA not only displays structural mimicry of WASP/WAVE family members, but also regulatory mimicry, having precisely co-opted the host machinery regulating these proteins. Comparisons based on ActA amino acid sequence suggest that unrelated pathogens that display actin-based motility may have evolved a similar strategy of regulatory mimicry.

## INTRODUCTION

The ability of various intracellular pathogens to spread from cell to cell without having to be exposed to the extracellular environment is a fundamental aspect of their pathogenic properties. To accomplish this important part of their infectious cycle, various bacterial pathogens have mastered the art of manipulating the host cell cytoskeleton (Gouin et al., 2005; Stevens et al., 2006). *Listeria monocytogenes* is a Gram-positive bacterium that displays the ability to invade mammalian epithelial cells and propagate into several epithelial structures, including the intestinal, fetoplacental and blood-brain barriers (Schlech, 1997). During intracytosolic growth, *L. monocytogenes* polymerizes host actin at one pole of the bacterial surface, allowing the bacteria to move through the cytosol via actin-based motility (Lambrechts et al., 2008). When the bacteria encounter the plasma membrane, the formation of membrane protrusions from an infected cell into an adjacent cell leads to the formation of a bacteria-containing double membrane vacuole in the

receiving cell. The bacteria escape this secondary vacuole and resume intracytosolic growth (Tilney and Portnoy, 1989).

*L. monocytogenes* actin-based motility relies on ActA (Brundage et al., 1993; Domann et al., 1992; Kocks et al., 1992), a bacterial factor that mimics the nucleation-promoting activity of WASP/WAVE family members (Goley and Welch, 2006; Pollard, 2007; Takenawa and Suetsugu, 2007) by recruiting the ARP2/3 complex to the bacterial surface (Welch et al., 1997, 1998).

The nucleation-promoting activity of WASP/WAVE family members relies on their VCA domain (V, verprolin) region (or WH2 domain), C (central) region, and the A (acidic) region. The VCA domain promotes nucleation by bringing together the first G-actin of a new filament and the ARP2/3 complex. The V region interacts with G-actin, and the C and A regions cooperate to recruit the ARP2/3 complex (Machesky and Insall, 1998; Marchand et al., 2001; Rohatgi et al., 1999). The C region is characterized by the presence of basic and hydrophobic residues that are essential for interaction with the ARP2/3 complex (Marchand et al., 2001; Panchal et al., 2003). In agreement with the importance of these basic residues in the C region of WASP, the R477K and K476E mutations (Fillat et al., 2001; Kolluri et al., 1995) have been identified in patients displaying the Wiskott-Aldrich syndrome (WAS), a form of immunodeficiency originally linked to mutations in the N-terminal part of WASP (Derry et al., 1994). The A region is characterized by the presence of a tryptophan residue embedded in a stretch of acidic residues. The exact function of this region is still unclear, but the tryptophan residue is essential for interaction with the ARP2/3 complex (Marchand et al., 2001).

Deletion analysis of ActA showed that the structural motif essential for interaction with the ARP2/3 complex in vitro and actin tail formation in vivo is a short stretch of basic residues located in the N-terminal part of the protein (Lasa et al., 1997; Skoble et al., 2000; Zalevsky et al., 2001), a region that displays similarities with the C region of WASP/WAVE VCA domain, including a stretch of basic residues (Bi and Zigmond, 1999).

In agreement with the importance of the conserved basic residues, introduction of the corresponding mutation in the N-terminal region of ActA (amino acid position 146–150) completely abolished actin tail formation as well as the recruitment of the ARP2/3 complex at the bacterial surface in vivo (Pistor et al., 2000). ActA does not display a region equivalent to the A region of WASP/WAVE in the vicinity of the conserved basic residues (C region). Instead, mapping and mutational analysis identified an acidic region harboring a key tryptophan residue located in the N-terminal domain of ActA (amino acid


position 41–45) (Boujemaa-Paterski et al., 2001; Skoble et al., 2000). Thus, similar to WASP and WAVE, two regions of ActA, displaying similarities with the C and A regions of the VCA domain of WASP and WAVE, cooperate for binding the ARP2/3 complex.

The nucleation-promoting activity of WASP/WAVE family members is tightly regulated in cells. For instance, all WASP family members harbor an autoinhibitory GBD domain that binds either to the C region of the VCA domain or to the Rho family GTPases (Goley and Welch, 2006; Pollard, 2007). WAVE family members do not display the autoinhibitory GBD domain. However, their nucleation-promoting activity is also tightly regulated through interaction with binding partners and posttranslational modifications (Takenawa and Suetsugu, 2007). Interestingly, recent studies have shown that the VCA domain of both WASP and WAVE 2 is subject to CK2-mediated phosphorylation, which increases their affinity for the ARP2/3 complex (Cory et al., 2003; Pocha and Cory, 2009). Although ActA was shown to be necessary and sufficient to promote the nucleation activity of the ARP2/3 complex in vitro (Welch et al., 1997), it is unclear whether host cell processes may be required to regulate the activity of ActA in vivo. Interestingly, ActA was shown to be highly phosphorylated upon infection in mammalian cells (Brundage et al., 1993), presumably by host cell kinase(s). However, the functional relevance of ActA phosphorylation in infected cells is unclear. Here, we show that, similar to WASP and WAVE, the affinity of ActA for the ARP2/3 complex is regulated by CK2-mediated phosphorylation. We discuss the implications of our findings with respect to the general mechanisms regulating the activity of nucleation-promoting factors and the evolution of functional mimicry by pathogens that hijack host cell cytoskeleton components.

## RESULTS

### CK2 Is Required for *L. monocytogenes* Spread from Cell to Cell

We recently conducted an RNAi screen for host factors involved in *L. monocytogenes* spread from cell to cell (R.S., unpublished data) and identified *CSNK2B* as a strong candidate gene. *CSNK2B* encodes the regulatory subunit of casein kinase 2 (CK2), a pleiotropic serine/threonine kinase catalyzing the phosphorylation of more than 300 substrates in eukaryotic cells (Pinna, 2002). To confirm the involvement of *CSNK2B* in *L. monocytogenes* spread from cell to cell, we used a pool of four redundant siRNA duplexes and silenced the expression of *CSNK2B* (see Experimental Procedures). In mock-treated cells, GFP-expressing *L. monocytogenes* formed a characteristic spreading pattern across the monolayer of cells (Figure 1A, mock-treated). In stark contrast, the bacteria accumulated in the center of a given infection foci in CK2-depleted cells, suggesting that the vast majority of the bacteria failed to spread to the neighboring cells (Figure 1A, *CSNK2B*-depleted). To quantify these phenotypes, we used computer-assisted image analysis to determine the spreading index (SI) as the number of cells harboring 75% of the total number of bacteria in a given infection foci (Figure S1 available online). When wild-type cells were infected for 8 hr, the spreading index was  $9 \pm 2$  (Figure 1A, mock-treated, image analysis SI = 9). In agreement with a role



**Figure 1. CK2 Is Required for *L. monocytogenes* Spread from Cell to Cell**

(A) Representative images of HeLa 229 cells (nuclei, blue channel) infected with GFP-expressing *L. monocytogenes* (Listeria, green channel) in mock-treated and CK2-depleted cells. The spreading index (SI) values and  $\pm$  SD were determined as described in Figure S1 by analyzing at least 12 foci of infection for each condition tested in three independent experiments.

(B and C) Efficiency of the silencing reagents. Cells were transfected with individual *CSNK2B* siRNA duplexes (A, B, C, and D) or a pool of the four silencing reagents (MIX) and were compared to mock-treated cells (CTRL) for silencing efficiency at the transcript level (B) and at the protein level (C). Values represent the mean  $\pm$  SD of three independent experiments.

(D) Cell-to-cell spreading phenotype in CK2-depleted cells using individual siRNA duplexes as described in (B) and (C) and quantified as shown in (A).

for *CSNK2B* in *L. monocytogenes* spread, the spreading index dropped to  $3 \pm 1$  in *CSNK2B*-depleted cells (Figure 1A, *CSNK2B*-depleted, image analysis, SI = 3).

To further validate the specific involvement of *CSNK2B*, we tested the four redundant silencing reagents individually. We confirmed their ability to silence *CSNK2B* expression at the transcript and protein levels (Figures 1B and 1C). We also determined their corresponding spreading index and confirmed their ability to confer the cell-to-cell spreading defect (Figure 1D). These results ruled out the possibility of an off-target effect and established that the observed cell-to-cell spreading phenotype is specific to *CSNK2B* depletion.

CK2 is a tetrameric enzyme displaying two redundant catalytic subunits, *CSNK2A1* and *CSNK2A2*, and two regulatory subunits, *CSNK2B*. We next tested whether *CK2 $\alpha$*  and/or *CK2 $\alpha'$*  may be involved in *L. monocytogenes* cell-to-cell spread. siRNA treatment targeting either *CSNK2A1* or *CSNK2A2* did not result in a significant spreading defect (Figure S2). However,

the simultaneous silencing of both *CSNK2A1* and *CSNK2A2* expression conferred a spreading defect phenotype similar to the one observed upon *CSNK2B* silencing (Figure S2). These results indicate that the CK2 holoenzyme is involved in *L. monocytogenes* cell-to-cell spread and that *CSNK2A1* and *CSNK2A2* isoforms probably play a (redundant) role as catalytic subunits.

### CK2 Is Required for Actin Tail Formation


To further analyze the role of CK2 in *L. monocytogenes* spread, we analyzed the formation of actin tails in mock-treated and CK2-depleted cells (Figure 2A). We scored bacteria that displayed an actin tail (Figure 2B, long and short tails) or were surrounded by F-actin (Figure 2B, clouds) or were not associated with F-actin at all (Figure 2B, no tails). In mock-treated cells, more than 90% of the bacteria were found in association with polymerized actin (Figure 2A, CTRL, and Figure 2C, CTRL). In CK2-depleted cells, the bacteria grew as microcolonies next to the nucleus (Figure 2A), and more than 70% of the bacteria were not associated with polymerized actin (Figure 2C). The rare bacteria that were associated with polymerized actin displayed very short tails or clouds and were often located next to the plasma membrane. Similar results were obtained in various cells types (Table S1). Altogether, these results indicate that the CK2 kinase is involved in *L. monocytogenes* cell-to-cell spread through the regulation of actin tail formation.

### CK2 Is Required for Recruitment of the ARP2/3 Complex

Two main host actin cytoskeleton regulators are involved in ActA-mediated actin tail formation. VASP binds a domain in ActA displaying proline-rich regions and enhances the actin tail elongation process (Lasa et al., 1995; Niebuhr et al., 1997; Smith et al., 1996). The ARP2/3 complex nucleates actin polymerization at the bacterial cell surface (Welch et al., 1998) and binds a domain in ActA displaying limited similarities to the C region of the VCA domain of WASP (Pistor et al., 2000; Skoble et al., 2000; Zalevsky et al., 2001). We first tested whether the actin tail formation defect observed in CK2-depleted cells may be related to a defect in VASP recruitment. In control cells, VASP was recruited to the cell surface of more than 95% of the bacteria and displayed a polar localization (Figure 2D, CTRL). In CK2-depleted cells, VASP recruitment was indistinguishable from the situation observed in control cells (Figure 2D, CSNK2B). These observations suggest that CK2 does not play a major role in VASP recruitment. We next tested the role of CK2 in the recruitment of the ARP2/3 complex. In control cells, the ARP2/3 complex colocalized with bacteria, as well as the actin tails (Figure 2E, CTRL). With some rare exceptions, the ARP2/3 complex did not colocalize with the bacteria in CK2-depleted cells (Figure 2E, CSNK2B). These data indicate that CK2 is required for efficient recruitment of the ARP2/3 complex to the bacterial cell surface in vivo.

### The C Region of ActA Is Phosphorylated by CK2 In Vitro

WASP/WAVE family members interact with the ARP2/3 complex through the C and A regions of their VCA domain (Machesky and Insall, 1998; Rohatgi et al., 1999). It was suggested that the N-terminal domain of ActA displays a stretch of basic residues (146-KKRRK-150) similar to the C region of WASP and WAVE (Bi and Zigmond, 1999) (see Figure 3A, C region).



**Figure 2. CK2 Is Required for Actin Tail Formation and Recruitment of the ARP2/3 Complex**


(A) Representative images of mock-treated (CTRL) and CSNK2B-depleted (CSNK2B) Swiss 3T3 cells infected with wild-type *L. monocytogenes* (green) and stained for DNA (blue) and F-actin (red).

(B) Scored actin tail patterns. The left images show F-actin only, and the right images show F-actin (red) and the bacteria (green).

(C) Histogram illustrating the quantification of scored actin tail patterns in mock-treated (CTRL) and CSNK2B-depleted (CSNK2B) cells. Values represent the mean  $\pm$  SD of three independent experiments.

(D and E) Images illustrating the recruitment of VASP (D) and ARP3 (E) in mock-treated (CTRL) and CK2-depleted (CSNK2B) Swiss 3T3 cells. For each image, the left inset shows the corresponding immunostaining only (VASP or ARP3), and the right inset shows immunostaining (red) and bacteria (green). The histograms on the left illustrate the quantification of the scored recruitment. Values represent the mean  $\pm$  SD of three independent experiments.

Structure/function analyses demonstrated that this short stretch of basic residues in ActA is essential for binding the ARP2/3 complex in vitro and for actin tail formation in vivo (Lasa et al., 1997; Pistor et al., 2000; Skoble et al., 2000). In addition,



**Figure 3. CK2 Phosphorylates the ARP2/3 Complex Binding Domain of ActA**

(A) Alignment of the peptide sequence of WASP, WAVE 2, ActA (ActA<sup>WT</sup>), and ActA mutated versions (ActA<sup>S155A,S157A</sup> and ActA<sup>S155E,S157E</sup>) showing the C region, the identified CK2 phosphorylation motifs, and the A region (position 41–45 in ActA) with the conserved W residue.

(B) Kinase assay with purified CK2 and GST fusion proteins displaying the wild-type C region of ActA (GST-ActA) or the S155A-S157A mutated version (GST-ActA<sup>S155A,S157A</sup>).

(C) MS/MS analysis of peptide spanning residues 151–170. The arrows indicate the MS/MS doubly charged y fragment ions for residues 153–156 (ASS<sup>\*</sup>D). The y14<sup>+2</sup> to the y18<sup>+2</sup> ions plus the y16-H<sub>3</sub>PO<sub>4</sub><sup>+2</sup> to the y18-H<sub>3</sub>PO<sub>4</sub><sup>+2</sup> ions demonstrate that Ser 155 is phosphorylated.

mapping and mutational analysis demonstrated the existence of a region in ActA equivalent to the A region of WASP/WAVE (Boujema-Paterski et al., 2001; Skoble et al., 2000) (see Figure 3A, A region). Interestingly, recent studies have shown that WASP and WAVE 2 are subject to CK2-mediated phosphorylation in the vicinity of their C region, which increases their affinity for the ARP2/3 complex (Cory et al., 2003; Pocha and Cory, 2009). Our comparative sequence analysis of WASP, WAVE 2, and ActA clearly indicated the presence of putative CK2 phosphorylation sites in the corresponding region of ActA (Figure 3A, SXXE/D). We used a kinase assay to show that the N-terminal region of ActA corresponding to amino acids 114–194 was phosphorylated by CK2 in vitro (Figure 3B, GST-ActA). The introduction of the S155A-S157A mutations in the corresponding N-terminal region of ActA (Figure 3A, ActA<sup>S155A,S157A</sup>) strongly decreased CK2-mediated phosphorylation, suggesting that residues S155 and S157 are potential targets for CK2 phosphorylation (Figure 3B, GST-ActA<sup>S155A,S157A</sup>).

### The C Region of ActA Is Phosphorylated In Vivo

We next used mass spectrometric approaches to specifically detect ActA phosphorylation in vivo. ActA was immunoprecipitated

from infected cell extracts and subjected to SDS electrophoresis (Brundage et al., 1993). The peptides resulting from in-gel digest of ActA and its phosphorylated products (Brundage et al., 1993) were fractionated into three pools and were analyzed by nano-LC tandem mass spectrometry for identification (Table S2). The first analyzed fraction represented 20% of the total protein digest (Figure S3A). The remaining 80% was enriched for phosphopeptides with TiO<sub>2</sub> affinity matrix, and both the enriched (Figure S3B) and the flow-through fractions (Figure S3C) were analyzed by LC-MS/MS. The data showed the enrichment of phosphopeptide <sup>151</sup>AIASSDSELES<sup>170</sup> KPTK<sup>170</sup> (harboring Ser155 and Ser157) as revealed by the presence of the triply charged phosphorylated ion (m/z 744.68) (Figures S3A and S3B) and the absence of the unmodified triply charged form of the peptide ion (m/z 718.03) (Figures S3B and S3C). The analysis of MS/MS spectrum revealed the presence of the doubly charged y fragment ions for residues 153–156 (Figure 3C). The y14<sup>+2</sup> to the y18<sup>+2</sup> ions and the y16-H<sub>3</sub>PO<sub>4</sub><sup>+2</sup> to the y18-H<sub>3</sub>PO<sub>4</sub><sup>+2</sup> ions demonstrated that Ser155 is phosphorylated (Figure 3C). We also conducted MS analysis in CK2-depleted cells for the nonphosphorylated version of the phosphopeptide of interest (m/z 718.0) and two (multiply charged)


versions of the detected phosphopeptide of interest (m/z 744.7, 1116.5) containing S155 and S157. The data show a consistent decrease for the signal corresponding to the phosphorylated AIASSDSELESLYDPKPTK peptide ions (Table S3). These results suggest that the relative level of phosphorylation of ActA on S155 is reduced in CK2-depleted cells.

### ActA Phosphorylation Regulates the Actin Tail Formation Process

To test the functional relevance of S155 and S157 phosphorylation upon infection, we introduced the single S155A and double S155A, S157A mutations by allelic exchange at the *actA* locus in the *L. monocytogenes* chromosome (see Experimental Procedures). The mutant bacteria displaying the *actA*<sup>S155A</sup> allele displayed no striking actin tail formation defect (data not shown). However, and similar to the situation observed in CSNK2B-depleted cells infected with wild-type bacteria, infection of mock-treated cells with mutant bacteria displaying the *actA*<sup>S155A, S157A</sup> allele revealed a strong spreading defect (Figure 4A, spreading index = 4 ± 2, n = 12), as well as a strong defect in actin tail formation (Figures 4A and 4C, *actA*<sup>WT</sup> and *actA*<sup>S155A, S157A</sup>). The *actA*<sup>S155A, S157A</sup> mutant bacteria were as infectious as wild-type, and the expression level of ActA<sup>S155A, S157A</sup> was similar to the expression level of ActA<sup>WT</sup> in infected cells (Figure S4). Moreover, the mutant bacteria expressing ActA<sup>S155A, S157A</sup> recruited VASP as efficiently as wild-type bacteria upon infection (Figure S5), indicating that ActA<sup>S155A, S157A</sup> was properly expressed at the bacterial cell surface. These results strongly suggested that phosphorylation on S155 and S157 is required for actin tail formation. In agreement with this assumption, we also showed that the introduction of the phospho-mimic S155E-S157E mutations in *actA* could partially rescue the actin polymerization defect observed in CK2-depleted cells (Figures 4B and 4C, *actA*<sup>WT</sup> and *actA*<sup>S155E, S157E</sup>). Similar to ActA<sup>S155A, S157A</sup>, ActA<sup>S155E, S157E</sup> was properly expressed at the bacterial cell surface (Figures S4 and S5). We noted that the vast majority of the mutant bacteria was associated with massive accumulation of F-actin, forming clouds or short tails, but failed to form long tails (Figures 4B and 4C, *actA*<sup>S155E, S157E</sup>), a defect also observed in wild-type cells infected with the *actA*<sup>S155E, S157E</sup> mutant bacteria (data not shown). Despite the formation of actin clouds and short tails, the bacteria displaying the phospho-mimic S155E-S157E mutations in *actA* did not spread properly (Figure 4B, spreading index = 4 ± 2, n = 12). These data indicate that low level of phosphorylation, as mimicked by the S155A-S157A mutations, as well as high level of phosphorylation, as mimicked by the S155E-S157E mutations, result in actin tail formation defects, suggesting that the phosphorylation level of S155 and S157 must be tightly regulated in vivo in order to achieve wild-type tail formation and efficient cell-to-cell spread.

### CK2 Phosphorylation Regulates Interaction of ActA with the ARP2/3 Complex


We next tested whether the actin tail formation defects observed with the mutant bacteria expressing ActA<sup>S155A, S157A</sup> may be related to a defect in the recruitment of the ARP2/3 complex. In mock-treated cells, the ARP2/3 complex colocalized with bacteria, as well as the actin tails (Figure 5A, *actA*<sup>WT</sup>). The



**Figure 4. ActA Phosphorylation Regulates Actin Tail Formation and Cell-to-Cell Spread**


(A and B) Cells infected with GFP-expressing bacteria (green) and stained for DNA (blue) and F-actin (red). The high-magnification images displayed at the bottom of each image show F-actin only (left) and F-actin (red) and bacteria (green) (right). The corresponding spreading index values and ± SD were determined by image analysis of at least 12 foci of infection in three independent experiments, as described in Figure S1 (*actA*<sup>WT</sup> in mock-treated cells, SI = 9 ± 3; *actA*<sup>S155A-S157A</sup> in mock-treated cells, SI = 4 ± 2; *actA*<sup>WT</sup> in CK2-depleted cells, SI = 3 ± 1; and *actA*<sup>S155E-S157E</sup> in CK2-depleted cells, SI = 3 ± 1).

(C) Histogram scoring actin tail formation as presented in (A) and (B). Values represent the mean ± SD of three independent experiments.



**Figure 5. ActA Phosphorylation Regulates ARP2/3 Complex Binding** (A and B) Mock-treated and CK2-depleted cells infected with GFP-expressing bacteria (green) and stained for DNA (blue) and ARP3 (red). The high-magnification images displayed at the bottom of each image show ARP3 only (left) and ARP3 (red) and bacteria (green) (right). The histograms on the right illustrate the quantification of the scored recruitment. Values represent the mean  $\pm$  SD of three independent experiments. (C) Interaction of wild-type (WT) and mutant (S155A-S157A and S155E-S157E) GST fusion proteins with purified ARP2/3 complex and western blot analysis (left) and quantification of three independent experiments (right). ARP3 signals were normalized with the GST values, and the normalized signal for WT was given an arbitrary value of 1 (relative intensity). Values represent the mean  $\pm$  SD of three independent experiments.

ARP2/3 complex did not colocalize with the bacteria expressing ActA<sup>S155A, S157A</sup> (Figure 5A, actA<sup>S155A, S157A</sup>), indicating that CK2 phosphorylation on S155-S157 is required for efficient recruitment of the ARP2/3 complex to the bacterial cell surface in vivo. Accordingly, the introduction of the phospho-mimic S155E-S157E mutations in actA could partially rescue the ARP2/3 complex recruitment defect observed in the CK2-depleted cells infected with wild-type bacteria (Figure 5B, actA<sup>WT</sup> and actA<sup>S155E, S157E</sup>). We next tested in vitro whether CK2-mediated phosphorylation on S155-S157 may affect the affinity of ActA for the ARP2/3 complex. To this end, we purified the N-terminal domain of ActA (1–194) as a GST fusion protein



**Figure 6. ActA Phosphorylation Affects *L. monocytogenes* Infection** Data showing colony-forming units (cfu) harvested from the spleen of three mice infected with wild-type (WT), ActA mutant ( $\Delta$ ActA), actA<sup>S155A, S157A</sup>, and actA<sup>S155E, S157E</sup> strains. Two-tailed Student's t tests were performed. The experiment was repeated twice with similar results.


and used a pull-down assay to analyze the interaction of ActA with purified ARP2/3 complex. In vitro phosphorylation of GST-ActA<sup>WT</sup> with purified CK2 dramatically increased the interaction with the ARP2/3 complex (Figure 5C, GST-ActA<sup>WT</sup> and GST-ActA<sup>WT</sup> + CK2). In agreement with our in vivo observations, the introduction of the S155A-S157A mutation completely abolished the effect of the CK2-mediated phosphorylation (Figure 5C, GST-ActA<sup>S155A-S157A</sup> and GST-ActA<sup>S155A-S157A</sup> + CK2). Also in agreement with our in vivo observations, the introduction of the phospho-mimic S155E-S157E mutations also increased the interaction of ActA with the ARP2/3 complex (Figure 5C, GST-ActA<sup>S155E-S157E</sup>), although the increase was not as dramatic as the one conferred by phosphorylation on S155 and S157 (8-fold versus 47-fold) (Figure 5C). Altogether, our in vivo and in vitro analyses demonstrate that CK2-mediated phosphorylation of ActA on S155 and S157 regulates its interaction with the ARP2/3 complex.

**ActA Phosphorylation Affects *Listeria monocytogenes* Infection**

We further tested the importance of the S155-S157 phosphorylation of ActA in a mouse model of infection. As a measure of the infection process, we found that the ability of the strains harboring the S155A-S157A or S155E-S157E actA alleles to colonize the spleen was clearly attenuated, displaying an intermediate phenotype, as compared to spleen colonization by the wild-type strain and the actA deletion mutant strain (Figure 6). We conclude that the regulation of actin tail formation through the CK2-mediated phosphorylation of ActA constitutes an important aspect of *L. monocytogenes* infection.

**Conservation of CK2 Phosphorylation Motifs in Nucleation-Promoting Factors**

Structural analysis of the C region of WASP revealed the presence of conserved hydrophobic and charged residues



**Figure 7. Peptide Sequence Alignment Showing the Presence of CK2 Phosphorylation Motifs in the Vicinity of the C Region of WASP/WAVE Family Members**

The conserved hydrophobic and arginine residues in the C region are boxed in green and red, respectively. The red dots indicate putative CK2 phosphorylation sites (pS/TXXD/E).

predicted to form an amphipathic helix interacting with the ARP2/3 complex and the WASP autoinhibitory domain (Panchal et al., 2003). We conducted comparative sequence analysis showing that the C region of all WASP/WAVE family members examined, including WASP-like factors produced by bacterial and viral pathogens such as *Listeria spp.*, *Rickettsia spp.*, *Wolbachia spp.*, and *Baculoviruses*, displays a signature that includes a perfectly conserved arginine residue surrounded by two hydrophobic residues (Figure 7). Importantly, with the exception of WASH (Linardopoulou et al., 2007), all WASP/WAVE-like family members examined also displayed putative

CK2 phosphorylation sites (S/TXXD/E) in the vicinity of the C region (Figure 7). Our analysis therefore supports the notion that, as shown for ActA in this study, unrelated virulence factors displaying nucleation-promoting activity, such as RickA (Gouin et al., 2004; Jeng et al., 2004) and p78/83 (Goley et al., 2006), may also be regulated by CK2-mediated phosphorylation (Figure 7). ActA, and potentially RickA and p78/83, constitutes a striking example of virulence factors that not only evolved as a functional mimicry of a host cell activity, but also precisely co-opted the host cell mechanism modulating this activity.

## DISCUSSION

In this study, we demonstrate that a host cell serine/threonine kinase, CK2, is required for efficient cell-to-cell spread of the intracellular pathogen *L. monocytogenes*. We show that the CK2-mediated phosphorylation of the ARP2/3-binding domain of ActA is required for proper actin tail formation and regulates the affinity of ActA for the ARP2/3 complex. These findings have implications for our general understanding of the mechanisms regulating host cell activities and the evolution of microbial factors mimicking these host cell activities.

Host cell nucleation-promoting factors of the WASP/WAVE family bind and activate the ARP2/3 complex through the C and A regions of their VCA domain (Goley and Welch, 2006; Pollard, 2007). Biochemical studies have suggested that the C region of WASP, N-WASP, and WAVE harbors a conserved arginine residue (position 477 in WASP) (Figure 6) and several conserved hydrophobic residues folding into an amphipathic helix that constitute the structural motif interacting with the autoinhibitory GBD domain, as well as the ARP2/3 complex (Panchal et al., 2003). Our comparative analyses of the C region of WASP/WAVE family members and the corresponding C region found in nucleation-promoting factors expressed by intracellular microbial pathogens revealed the strict conservation of the arginine residue and two surrounding hydrophobic residues (Figure 7). This conservation is, however, less apparent for the newly identified WASP-like factors, including WHAMM (Campellone et al., 2008) and JMY (Zuchero et al., 2009) (Figure 7). We propose that this signature constitutes the minimal structural motif required for interaction with the ARP2/3 complex with the C region and, as such, constitutes the structural basis for the functional mimicry displayed by the C region of ActA (Figure 7).

We present genetic and biochemical evidence showing that phosphorylation of the C region of ActA on S155 and S157 is important for the recruitment of the ARP2/3 complex. Both sites qualify as bona fide CK2 phosphorylation sites (consensus sequence SXXD/E). We also present compelling evidence suggesting that S155 is a direct target for CK2 phosphorylation in vitro and in vivo. However, we were not able to detect S157 phosphorylation in vivo. Thus, although very likely, the actual involvement of CK2 in S157 phosphorylation remains to be demonstrated.

Similar to ActA, the CK2-mediated phosphorylation of the C region of WASP and WAVE 2 (see Figure 3A) was shown to increase the affinity of the VCA domain for the ARP2/3 complex (Cory et al., 2003; Pocha and Cory, 2009). It is noteworthy that the vast majority of CK2 phosphorylation sites are serine/threonine residues surrounded by acidic residues from position -1 to +4, with an absolute conservation in position +3 (S/TXXE/D) (Pinna, 2002). Interestingly, the acidic residues surrounding S155 and S157 were identified as important for ActA activity in a systematic charged-to-alanine mutational analysis of the N-terminal region of ActA (see Lauer et al., 2001; mutant 48, 155-SDSELE > 155-SASALA). Taking all of these observations together, we propose a model in which the negative charges provided by the acidic residues surrounding the phosphorylation sites contribute to the affinity of the C region for the ARP2/3 complex (Figure 3A). In this model, the addition and

removal of phosphate groups modulate the interaction with the ARP2/3 complex. Accordingly, the in vitro phosphorylation of ActA on S155 and S157 dramatically increased the affinity of ActA for the ARP2/3 complex (Figure 5C). Moreover, the introduction of the S155A-S157A mutations in ActA led to a strong defect in actin tail formation and recruitment of the ARP2/3 complex in vivo (Figure 5A). Furthermore, the introduction of the phospho-mimic mutation S155E-S157E increased the affinity of ActA for the ARP2/3 complex in vitro (Figure 5C), and the corresponding mutant strain could recruit the ARP2/3 complex in CK2-depleted cells (Figure 5B). However, the affinity of the phosphorylated wild-type protein was significantly higher than the affinity of the phospho-mimic mutant. In addition, the rescue of actin tail formation in CK2-depleted cells was only partial, as the bacteria failed to form long tails (Figure 4C). These results may indicate that, in addition to the negative charges, the phosphate groups may provide additional features that contribute to the affinity of the C region for the ARP2/3 complex. It is also possible that, in vivo, in order to achieve proper actin tail formation, the level of ActA phosphorylation must be tightly regulated through phosphorylation/dephosphorylation cycles, perhaps suggesting the existence of a phosphatase activity that modulates the phosphorylation state of the C region in vivo. Thus, *L. monocytogenes* may provide a tractable system to understand the exact role of the CK2-mediated phosphorylation and to identify additional factors regulating the activity of the C region of WASP/WAVE family members.

With the exception of the conserved residues in the C region and putative CK2 phosphorylation sites, unrelated microbial factors displaying nucleation-promoting activities, such as ActA in *Listeria spp.*, RickA in *Rickettsia spp.* (Gouin et al., 2004; Jeng et al., 2004), and p78/83 in *Baculoviruses* (Goley et al., 2006; Machesky et al., 2001), do not display extensive amino acid sequence similarities with WASP/WAVE family members. Thus, it is likely that these microbial mimics of WASP and WAVE were generated by convergent evolution (Stebbins and Galan, 2001). In this context, it is remarkable that ActA, and potentially RickA and p78/83, co-opted a regulatory mechanism similar to WASP and WAVE. This probably indicates that the CK2-mediated phosphorylation represents a unique solution to the problem of regulating the cellular activity of nucleation-promoting factors displaying a C region. We propose that ActA defines a type of bacterial mimicry that we refer to as regulatory mimicry, whereby a pathogen has not only evolved a structural mimic of a host cell activity, such as the VCA domain and its role in the recruitment of the ARP2/3 complex, but has also precisely co-opted the host cell machinery that regulates this host cell activity, such as the serine/threonine kinase CK2 and its role in regulating the affinity of the VCA domain for the ARP2/3 complex.

## EXPERIMENTAL PROCEDURES

### Bacterial and Eukaryotic Cell Growth Conditions

*L. monocytogenes* strain 10403S was grown overnight in BHI (Difco) at 30°C without agitation prior to infection. HeLa 229 cells (ATCC #CCL-2.1) and Swiss 3T3 (ATCC #CCL 92) cells were grown in DMEM (Invitrogen) supplemented with 10% FBS (GIBCO) at 37°C in a 5% CO<sub>2</sub> incubator.

### Listeria monocytogenes Infection

Cells were infected (MOI = 20) with *Listeria monocytogenes* strain (10403S) expressing GFP under the control of an IPTG-inducible promoter (H.A., unpublished data). The plates were centrifuged at 1000 rpm for 5 min, and internalization of the bacteria was allowed to proceed for 1 hr at 37°C before gentamicin (50 μM final) and IPTG (4 mM final) were added in order to kill the remaining extracellular bacteria and visualize internalized bacteria. The cells were incubated at 37°C for 6–8 hr and then fixed in PBS 4% formaldehyde at 4°C overnight. The cells were stained with Hoechst (Molecular Probes, Inc.), washed in PBS, and kept at 4°C until imaging.

### CSNK2B Silencing and Validation Procedure

Cells were transfected by reverse transfection with Dharmafect1 and individual *CSNK2B* siRNA (A, B, C, and D, 50 nM final) or a pool of the four silencing reagents (12.5 nM each, 50 nM total) and were incubated for 72 hr on coverslips in a 24-well plate format. The sequences targeted by the individual siRNA duplexes were as follows: A, CAACCAGAGUGACCUGAUU; B, GACAAGCU CUAGACAUGAU; C, CAGCCGAGAUGCUUUUAUGG; D, GCUCUACGGUUUC AAGAUC.

For real-time PCR analysis, total RNA and first-strand cDNA synthesis was performed using the TaqMan Gene Expression Cells-to-Ct Kit (Applied Biosystems), as recommended by the manufacturer, with the addition of DNaseI for the removal of unwanted genomic DNA. *CSNK2B* mRNA levels were determined by quantitative real-time PCR using the LightCycler 480 SYBR Green 1 Master Kit and LightCycler 480 instrument (Roche Biochemicals, Indianapolis, IN).

For western blot analysis, cells were transfected with the corresponding silencing reagents in a 24-well plate format, and cells were lysed directly in Laemmli sample buffer. The anti-CSNK2B antibody (1:200) was obtained from Santa Cruz (sc-20710). The anti-actin antibody (1:10,000) was obtained from Sigma (A2066). The anti-rabbit HRP-conjugated secondary antibody (1:10,000) was obtained from Jackson ImmunoResearch (111-035-14).

### Immunofluorescence Procedures

Swiss 3T3 cells were seeded on glass coverslips, infected with *L. monocytogenes*, and fixed in PBS 5% paraformaldehyde. The cells were permeabilized and stained in PBS/0.05% Saponin/1% BSA containing the appropriate primary antibody for 2 hr at room temperature and were washed five times in PBS/0.05% Saponin/1% BSA before addition of secondary antibodies in PBS/0.05% Saponin/1% BSA for 1 hr at room temperature. The samples were then mounted onto glass slides using FluoroGuard Antifade Reagent (Bio-Rad). Rabbit monoclonal anti-VASP antibody (1:100) was obtained from Cell Signaling (Clone 9A2). Rabbit polyclonal anti-Arp3 antibody (1:100) was obtained from Upstate (07-272). Alexa-fluor 488 secondary antibody (1:1000), Alexa-fluor 568 conjugated phalloidin (1:500), and Hoechst (1:500) were obtained from Invitrogen.

### ActA Mutagenesis

The S155A-S157A double mutation (TCG > GCA and AGT > GCA, respectively) was introduced using the overlapping primers 5S155A-S157A (AGAA GAAAAGCCATAGCGTCGCGAGATGCGAGAGCTTGAAAGCCTTACTTAT) and 3S155A-S157A (ATAAGTAAGGCTTTCAAGCTCTGCATCTGCCAGCGCTATG GCTTTTCTTCT). The construct was generated by overlap PCR using the flanking 5' primer ActA5Eco (GAAGAATTCGCGCTTCAACAAGCAGCG) with 3S155A-S157A and 3' primer ActA3BamHI (GGAGGATCCCCTTATCAGA GCCGGCGCGC) with 5S155A-S157A. The two overlapping fragments were gel purified and used as template for PCR amplification using ActA5Eco and ActA3BamHI as flanking primers. The resulting 1.6 kb PCR fragment encoding the corresponding S155AS157A mutation was digested by EcoRI and BamHI and cloned into pMAD (Arnaud et al., 2004). The resulting plasmid, pMAD-ACTA<sup>S155AS157A</sup>, was introduced into *L. monocytogenes* 10403S by electroporation, and the gene replacement was conducted as previously described (Arnaud et al., 2004). The resulting mutant strain (10403S actA<sup>S155A-S157A</sup>) was verified by PCR amplification using chromosomal DNA of the corresponding strain and direct DNA sequencing of the PCR product. The actA<sup>S155E-S157E</sup> mutant strain was generated similarly by generating plasmid pMAD-ACTA<sup>S155ES157E</sup> using the primers 5S155E-S157E (AGAAGAAAAG CCATAGCGTCGGAAAGATGAAGAGCTTGAAAGCCTTACTTAT) and 3S155E-

S157E (ATAAGTAAGGCTTTCAAGCTCTTTCATCTTCCGACGCTATGGCTTTT CTTCT).

### In Vitro Kinase Assay

The DNA fragment corresponding to amino acids 114–194 of ActA was PCR amplified from chromosomal DNA using primers 5GST1Eco (GAAGAATTC TAGTGGCTATAAATGAAGAGGCTTCA) and 3GSTH3 (AAGAAGCTTCTGC ATGCTAGAATCTAAGTCACTTTC), digested with EcoRI and HindIII, and cloned into pGEX-KG (Amersham Pharmacia Biotech). The S155A-S157A double mutation was amplified using 5GST1Eco and 3GSTH3 as primers and pMAD-ACTA<sup>S155AS157A</sup> as template. Both constructs were verified by DNA sequencing. The corresponding fusion proteins were produced in *E. coli* BL21 (Invitrogen). The bacteria were grown at 30°C until OD<sub>600</sub> = 0.5, and expression was induced by adding 0.5 mM IPTG for 1 hr. After resuspension of the bacteria in PBS and disruption of the bacterial cell wall by sonication, the fusion proteins were purified using glutathione Sepharose 4B (GS4B) (Amersham Pharmacia Biotech). To conduct the kinase assay, fusion proteins attached to the GS4B beads were washed and resuspended in kinase buffer (200 mM Tris-HCl [pH 7.5], 10 mM KCl, 10 mM MgCl<sub>2</sub>, 100 μM ATP, and 200 μM sodium orthovanadate) supplemented with 10 μCi (<sup>γ</sup>-<sup>32</sup>P)ATP and 50 U CK2 (New England Biolabs, Inc) and incubated for 10 min at 30°C. GS4B beads were washed three times in PBS, resuspended in 30 μl Laemmli buffer (Sigma), and subjected to SDS-PAGE analysis. The gel was stained with Coomassie brilliant blue and dried, and incorporated <sup>32</sup>P was detected using autoradiography.

### Mass Spectrometric Analysis

#### In-Gel Protein Digestion

Coomassie stained gel bands corresponding to ActA protein isolated in one-dimensional polyacrylamide gel were subjected to in situ enzymatic digestion. The gel plugs were washed with 250 μl 50% acetonitrile/50% water for 5 min followed by 250 μl of 50 mM ammonium bicarbonate/50% acetonitrile/50% water for 30 min. One final wash was performed using 10 mM ammonium bicarbonate/50% acetonitrile/50% water for 30 min. After washing, the gel plugs were dried in a Speedvac and rehydrated with 0.1 μg of modified trypsin (Promega) per ~15 mm<sup>3</sup> of gel in 15 μl 10 mM ammonium bicarbonate. Samples were digested at 37°C for 16 hr.

#### Titanium Dioxide Enrichment

The digest was acidified with 0.5% TFA and 50% acetonitrile. Top Tips (Glyen Corp.) were prepared by washing three times with 40 μl of each 100% acetonitrile, followed by 0.2 M sodium phosphate (pH 7.0) and 0.5% TFA and 50% acetonitrile. Washes were spun through into an Eppendorf tube at 2000 rpm for 1 min. The acidified digest supernatant was loaded into the Top Tip, spun at 1000 rpm for 1 min, and then spun at 3000 rpm for 2 min. Gel pieces were rinsed with 40 μl 0.5% TFA and 50% acetonitrile, with the supernatant transferred to the Top Tip and the spin repeated. The Top Tip was then washed with 40 μl 0.5% TFA and 50% acetonitrile, and the spin was repeated. The flowthrough from these washes was saved and analyzed by LC-MS/MS as below. Phosphopeptides were eluted from the Top Tip with 30 μl 28% ammonium hydroxide (three times). Both the flowthrough and eluted fractions were dried in Speedvac and redried in 40 μl water. Samples were dissolved in 3 μl 70% formic acid, vortexed, diluted with 7 μl 0.1% TFA, spun, and transferred to LC-MS/MS vials, where 5 μl was injected.

#### LC-MS/MS

The tandem mass spectral analysis was performed with a Thermo Fischer Scientific LTQ-Orbitrap mass spectrometer equipped with a Waters nano-Acquity UPLC system. Peptides/phosphopeptides were loaded onto a Waters Symmetry C18 180 μm × 20 mm trap column and separated through a 1.7 μm, 75 μm × 250 mm nanoAcquity UPLC column (35°C) immediately connected to the electrospray ionization source. Trapping was performed at 15 μl/min, 99% buffer A (100% water, 0.1% formic acid) for 1 min. Peptide separation was performed at 300 nl/min with buffer A (100% water, 0.1% formic acid) and buffer B (100% CH<sub>3</sub>CN, 0.075% formic acid). A linear gradient (51 min) was run with 5% buffer B at initial conditions, 50% B at 50 min, and 85% B at 51 min. MS was acquired in the LTQ-Orbitrap using 1 microscan and a maximum inject time of 900 microseconds. A precursor mass spectral acquisition was taken in the orbitrap followed by four data-dependent MS/MS acquisitions in the ion trap. MS<sup>3</sup> scans were also obtained if neutral loss of a phosphate group was

detected corresponding to m/z differences of 98.0, 49.0, and 32.7. The data was searched using the Mascot (Matrix Science) database search algorithm against the NCBI nr protein database.

#### In Vitro Protein Binding Assay

Plasmids pGEX-GST3, pGEX-GST3<sup>S155AS157A</sup>, and pGEX-GST3<sup>S155ES157E</sup> were generated using oligonucleotides 5GST3 (GAAGAATTCGTGGG ATTAATTAGATTATGCGTGC) and 3GSTH3 (AAGAAGCTTCTGCATG CTAGAATCTAAGTCACTTTC) as primers and chromosomal DNA, pMAD-ACTA<sup>S155AS157A</sup>, and pMAD-ACTA<sup>S155ES157E</sup> as templates, respectively. The corresponding BL21 strains were grown at 37°C to an O.D. of 0.5 before addition of IPTG (1 mM final concentration) and further incubation for 1 hr at 30°C. Bacteria were subsequently harvested in 1 ml PBS supplemented with 10 μl of a protease inhibitor cocktail (Sigma) and lysed by sonication. The GST fusion proteins were immobilized on 200 μl of glutathione Sepharose beads (GE Healthcare) for 2 hr at 4°C. CK2-treated beads were incubated at 30°C for 20 min in CK2 reaction buffer (20 mM Tris-HCl [pH 7.5], 50 mM KCl, and 10 mM MgCl<sub>2</sub>) supplemented with 200 μM ATP and 50 U of purified CK2 enzyme (New England Biosciences). Untreated and CK2-treated beads were washed once in XB (10 mM HEPES [pH 7.6], 100 mM KCl, 1 mM MgCl, 0.1 mM EDTA, and 1 mM DTT) and subsequently incubated with 5 μM purified bovine Arp2/3 complex (Cytoskeleton) in XB plus 0.1% (v/v) Tween-20, 0.2 mM ATP, and 0.5 mg/ml BSA (50 nM) at 4°C for 2 hr. Beads were subsequently washed twice in 150 μl of XB plus 0.1% (v/v) Tween-20 and 0.2 mM ATP. Beads were resuspended in 40 μl of Laemmli sample buffer, boiled for 10 min, and subjected to western blot analysis. Membranes were incubated in blocking buffer (PBS plus 5% dry nonfat milk) and probed with either rabbit polyclonal Anti-Arp3 (1:500) obtained from Millipore or rabbit polyclonal Anti-GST (1:250) obtained from Santa Cruz Biotechnology. IRDye800-conjugated anti-rabbit IgG (Rockland, Gilbertsville, PA) were used as secondary antibodies. Signal intensities were quantified using the Odyssey infrared image system (Li-Cor).

#### Mouse Infection

Groups of four 8-week-old female BALB/c mice were infected intraperitoneally with  $5 \times 10^4$  bacteria of a given genotype. At 4 days postinfection, the animals were euthanized, spleens were collected and homogenized in PBS, and the number of colony-forming units (cfus) was determined on BHI agar plates.

#### SUPPLEMENTAL DATA

Supplemental Data include five figures and three tables and can be found with this article online at [http://www.cell.com/molecular-cell/supplemental/S1931-3128\(09\)00285-6](http://www.cell.com/molecular-cell/supplemental/S1931-3128(09)00285-6).

#### ACKNOWLEDGMENTS

We thank Dan Portnoy for the generous gift of the ActA antibody. We thank Jorge Galan for suggesting the mouse experiments. We thank Neal Gilksman for helping with image analysis. We thank Isabelle Derré, Ana-Maria Dragoi, Jorge Galan, and Craig Roy for critical reading of the manuscript. This work was supported by the Mallinckrodt foundation (H.A.), the National Institutes of Health training grant T32 AI-07640 (R.C.), and a Northeast Biodefense Center grant U54 AI-057158 for Proteomics Core (K.L.S. and E.E.G.) and grant R01 AI-073904 (H.A.).

Received: January 16, 2009

Revised: June 17, 2009

Accepted: August 25, 2009

Published: September 16, 2009

#### REFERENCES

Arnaud, M., Chastanet, A., and Debarbouille, M. (2004). New vector for efficient allelic replacement in naturally nontransformable, low-GC-content, gram-positive bacteria. *Appl. Environ. Microbiol.* 70, 6887–6891.

Bi, E., and Zigmond, S.H. (1999). Actin polymerization: Where the WASP stings. *Curr. Biol.* 9, R160–R163.

Boujemaa-Paterski, R., Gouin, E., Hansen, G., Samarin, S., Le Clairche, C., Didry, D., Dehoux, P., Cossart, P., Kocks, C., Carlier, M.F., et al. (2001). *Listeria* protein ActA mimics WASp family proteins: it activates filament barbed end branching by Arp2/3 complex. *Biochemistry* 40, 11390–11404.

Brundage, R.A., Smith, G.A., Camilli, A., Theriot, J.A., and Portnoy, D.A. (1993). Expression and phosphorylation of the *Listeria monocytogenes* ActA protein in mammalian cells. *Proc. Natl. Acad. Sci. USA* 90, 11890–11894.

Campellone, K.G., Webb, N.J., Znameroski, E.A., and Welch, M.D. (2008). WHAMM is an Arp2/3 complex activator that binds microtubules and functions in ER to Golgi transport. *Cell* 134, 148–161.

Cory, G.O., Cramer, R., Blanchoin, L., and Ridley, A.J. (2003). Phosphorylation of the WASP-VCA domain increases its affinity for the Arp2/3 complex and enhances actin polymerization by WASP. *Mol. Cell* 11, 1229–1239.

Derry, J.M., Ochs, H.D., and Francke, U. (1994). Isolation of a novel gene mutated in Wiskott-Aldrich syndrome. *Cell* 78, 635–644.

Domann, E., Wehland, J., Rohde, M., Pistor, S., Hartl, M., Goebel, W., Leimeister-Wachter, M., Wuenscher, M., and Chakraborty, T. (1992). A novel bacterial virulence gene in *Listeria monocytogenes* required for host cell microfilament interaction with homology to the proline-rich region of vinculin. *EMBO J.* 11, 1981–1990.

Fillat, C., Espanol, T., Oset, M., Ferrando, M., Estivill, X., and Volpini, V. (2001). Identification of WASP mutations in 14 Spanish families with Wiskott-Aldrich syndrome. *Am. J. Med. Genet.* 100, 116–121.

Goley, E.D., and Welch, M.D. (2006). The ARP2/3 complex: an actin nucleator comes of age. *Nat. Rev. Mol. Cell Biol.* 7, 713–726.

Goley, E.D., Ohkawa, T., Mancuso, J., Woodruff, J.B., D'Alessio, J.A., Cande, W.Z., Volkman, L.E., and Welch, M.D. (2006). Dynamic nuclear actin assembly by Arp2/3 complex and a baculovirus WASP-like protein. *Science* 314, 464–467.

Gouin, E., Egile, C., Dehoux, P., Villiers, V., Adams, J., Gertler, F., Li, R., and Cossart, P. (2004). The RickA protein of *Rickettsia conorii* activates the Arp2/3 complex. *Nature* 427, 457–461.

Gouin, E., Welch, M.D., and Cossart, P. (2005). Actin-based motility of intracellular pathogens. *Curr. Opin. Microbiol.* 8, 35–45.

Jeng, R.L., Goley, E.D., D'Alessio, J.A., Chaga, O.Y., Svitkina, T.M., Borisy, G.G., Heinzen, R.A., and Welch, M.D. (2004). A *Rickettsia* WASP-like protein activates the Arp2/3 complex and mediates actin-based motility. *Cell. Microbiol.* 6, 761–769.

Kocks, C., Gouin, E., Tabouret, M., Berche, P., Ohayon, H., and Cossart, P. (1992). *L. monocytogenes*-induced actin assembly requires the actA gene product, a surface protein. *Cell* 68, 521–531.

Kolluri, R., Shehabeldin, A., Peacocke, M., Lamhonwah, A.M., Teichert-Kuliszewska, K., Weissman, S.M., and Siminovitich, K.A. (1995). Identification of WASP mutations in patients with Wiskott-Aldrich syndrome and isolated thrombocytopenia reveals allelic heterogeneity at the WAS locus. *Hum. Mol. Genet.* 4, 1119–1126.

Lambrechts, A., Gevaert, K., Cossart, P., Vandekerckhove, J., and Van Troys, M. (2008). *Listeria comet tails*: the actin-based motility machinery at work. *Trends Cell Biol.* 18, 220–227.

Lasa, I., David, V., Gouin, E., Marchand, J.B., and Cossart, P. (1995). The amino-terminal part of ActA is critical for the actin-based motility of *Listeria monocytogenes*; the central proline-rich region acts as a stimulator. *Mol. Microbiol.* 18, 425–436.

Lasa, I., Gouin, E., Goethals, M., Vancompernelle, K., David, V., Vandekerckhove, J., and Cossart, P. (1997). Identification of two regions in the N-terminal domain of ActA involved in the actin comet tail formation by *Listeria monocytogenes*. *EMBO J.* 16, 1531–1540.

Lauer, P., Theriot, J.A., Skoble, J., Welch, M.D., and Portnoy, D.A. (2001). Systematic mutational analysis of the amino-terminal domain of the *Listeria monocytogenes* ActA protein reveals novel functions in actin-based motility. *Mol. Microbiol.* 42, 1163–1177.

- Linardopoulou, E.V., Parghi, S.S., Friedman, C., Osborn, G.E., Parkhurst, S.M., and Trask, B.J. (2007). Human subtelomeric WASH genes encode a new subclass of the WASP family. *PLoS Genet.* 3, e237.
- Machesky, L.M., and Insall, R.H. (1998). Scar1 and the related Wiskott-Aldrich syndrome protein, WASP, regulate the actin cytoskeleton through the Arp2/3 complex. *Curr. Biol.* 8, 1347–1356.
- Machesky, L.M., Insall, R.H., and Volkman, L.E. (2001). WASP homology sequences in baculoviruses. *Trends Cell Biol.* 11, 286–287.
- Marchand, J.B., Kaiser, D.A., Pollard, T.D., and Higgs, H.N. (2001). Interaction of WASP/Scar proteins with actin and vertebrate Arp2/3 complex. *Nat. Cell Biol.* 3, 76–82.
- Niebuhr, K., Ebel, F., Frank, R., Reinhard, M., Domann, E., Carl, U.D., Walter, U., Gertler, F.B., Wehland, J., and Chakraborty, T. (1997). A novel proline-rich motif present in ActA of *Listeria monocytogenes* and cytoskeletal proteins is the ligand for the EVH1 domain, a protein module present in the Ena/VASP family. *EMBO J.* 16, 5433–5444.
- Panchal, S.C., Kaiser, D.A., Torres, E., Pollard, T.D., and Rosen, M.K. (2003). A conserved amphipathic helix in WASP/Scar proteins is essential for activation of Arp2/3 complex. *Nat. Struct. Biol.* 10, 591–598.
- Pinna, L.A. (2002). Protein kinase CK2: a challenge to canons. *J. Cell Sci.* 115, 3873–3878.
- Pistor, S., Grobe, L., Sechi, A.S., Domann, E., Gerstel, B., Machesky, L.M., Chakraborty, T., and Wehland, J. (2000). Mutations of arginine residues within the 146-KKRRK-150 motif of the ActA protein of *Listeria monocytogenes* abolish intracellular motility by interfering with the recruitment of the Arp2/3 complex. *J. Cell Sci.* 113, 3277–3287.
- Pocha, S.M., and Cory, G.O. (2009). WAVE2 is regulated by multiple phosphorylation events within its VCA domain. *Cell Motil. Cytoskeleton* 66, 36–47.
- Pollard, T.D. (2007). Regulation of actin filament assembly by Arp2/3 complex and formins. *Annu. Rev. Biophys. Biomol. Struct.* 36, 451–477.
- Rohatgi, R., Ma, L., Miki, H., Lopez, M., Kirchhausen, T., Takenawa, T., and Kirschner, M.W. (1999). The interaction between N-WASP and the Arp2/3 complex links Cdc42-dependent signals to actin assembly. *Cell* 97, 221–231.
- Schlech, W.F., III. (1997). *Listeria gastroenteritis*—old syndrome, new pathogen. *N. Engl. J. Med.* 336, 130–132.
- Skoble, J., Portnoy, D.A., and Welch, M.D. (2000). Three regions within ActA promote Arp2/3 complex-mediated actin nucleation and *Listeria monocytogenes* motility. *J. Cell Biol.* 150, 527–538.
- Smith, G.A., Theriot, J.A., and Portnoy, D.A. (1996). The tandem repeat domain in the *Listeria monocytogenes* ActA protein controls the rate of actin-based motility, the percentage of moving bacteria, and the localization of vasodilator-stimulated phosphoprotein and profilin. *J. Cell Biol.* 135, 647–660.
- Stebbins, C.E., and Galan, J.E. (2001). Structural mimicry in bacterial virulence. *Nature* 412, 701–705.
- Stevens, J.M., Galyov, E.E., and Stevens, M.P. (2006). Actin-dependent movement of bacterial pathogens. *Nat. Rev. Microbiol.* 4, 91–101.
- Takenawa, T., and Suetsugu, S. (2007). The WASP-WAVE protein network: connecting the membrane to the cytoskeleton. *Nat. Rev. Mol. Cell Biol.* 8, 37–48.
- Tilney, L.G., and Portnoy, D.A. (1989). Actin filaments and the growth, movement, and spread of the intracellular bacterial parasite, *Listeria monocytogenes*. *J. Cell Biol.* 109, 1597–1608.
- Welch, M.D., Iwamatsu, A., and Mitchison, T.J. (1997). Actin polymerization is induced by Arp2/3 protein complex at the surface of *Listeria monocytogenes*. *Nature* 385, 265–269.
- Welch, M.D., Rosenblatt, J., Skoble, J., Portnoy, D.A., and Mitchison, T.J. (1998). Interaction of human Arp2/3 complex and the *Listeria monocytogenes* ActA protein in actin filament nucleation. *Science* 281, 105–108.
- Zalevsky, J., Grigorova, I., and Mullins, R.D. (2001). Activation of the Arp2/3 complex by the *Listeria acta* protein. Acta binds two actin monomers and three subunits of the Arp2/3 complex. *J. Biol. Chem.* 276, 3468–3475.
- Zuchero, J.B., Coutts, A.S., Quinlan, M.E., Thangue, N.B., and Mullins, R.D. (2009). p53-cofactor JMY is a multifunctional actin nucleation factor. *Nat. Cell Biol.* 11, 451–459.